



THE CASE REPORTS ORPHANAGE OBSTETRICS MENOPAUSE WITH BLEEDING 2 MONTHS AT PREMARY HEALTH CARE NGESREP

Susiyati¹ Siti Nurjanah² Erna Kusumawati³ Ariyani Lutfitasari⁴

^{1,2}Midwife Professional Education Department, Universitas Muhammadiyah Semarang

^{3,4}Midwifery Undergraduate Department, Universitas Muhammadiyah Semarang

Corresponding author : sitinurjanah@unimus.ac.id

ABSTRACT

The world health organization (who) in asia , in 2025 number of women have menopause would rise from 107 million people being 373 million people. Suds menopause experienced by many women almost all the world around 70 - 80 % European women , 60 % in America , 57 % in Malaysia , 18 % in China , 10 % in Japan and also 10 % in Indonesia. According to the central bureau of statistics bps 2013 () years increase in life expectancy (UHH) in 2000 in indonesia is 64,5 years 7,18 % with the elderly population .This figure rose to 69,43 in in 2010 with the elderly population 7,56 % and in a year with the 2011 69,65 % 7,58 elderly population. With increasing life expectancy, is an increased occurrence of disease in women. old especially Women experiencing menopause, good, early menopause pre menopausal, perimenopause and post menopause general to experience symptoms of the top (climacteric) and transition or intermediate. Bleeding pascamenopause are complaints who was often experienced by women pascamenopause.Bleeding pascamenopause happens to about 4-11 % whose woman is menopause.10 % cases caused by the ferocity of, namely. endometrial cancer The highest insidensi endometrial cancer found in women pascamenopause aged 60-79 years.Case studies by the observational descriptive with the approach the case study method research conducted by means of research the problem through a case consisting of a single unit .This case study using the obstetrics care 7 varney from collecting a baseline data to evaluation and the development of using SOAP. Found the bleeding leading to pathology so done the act of collaboration with dr. SpOG for further examination . Examination laborat support , ultrasound and referred to the hospital. A woman can be said in menopause when the menstrual cycle stop for at least 12 months consecutive. After menopause, some women can still bleeding from the vagina called also as pascamenopause bleeding

Keywords : Menopause, pascamenopause bleeding

1. INTRODUCTION

Menopause is occurring in the natural every woman, and it is affected the different cultures and individual perception. To people in general, adult age having high appreciation than age esp women who had menopause.Menopause is a phenomenon in the life of women characterized by cessation the menstrual cycle within a the past years is undergo menstruation (Setyowati, 2018). The average age woman had menopause is 51 years, but can happen to age 40-45 years old and still normal category (Holland, 2015). In the before menopause, various complaints that will be experienced in women, menopause as irritable, fear, agitated, irritable, conflict heat (hot flushes), depression, headache, fast tired, difficult to concentrate, easily forgotten, less power, weight in, bone pain and muscles, sleep disorders, obstipasi, the beating heart.

The world health organization (WHO) in Asia, in 2025 the number of women menopause will increase from 107 million souls into 373 million people .Suds menopause for many women almost all the world around 70-80 % European women , 60 % in America , 57 % in Malaysia , 18 % in China , 10 % in Japan and also 10 % (Fitriani 2018). Based on the calculation of statistics in 2020 country s population is expected to reach 262,6 million inhabitants and the number of women who live in of your at between 45-55 years is about 30,3 ruswanti million people) 2018. According to the central bureau of statistics bps 2013 years increase in life expectancy (UHH) in 2000 in indonesia is 64,5 years 7,18 % with the elderly population .This figure rose to 69,43 in in 2010 with the elderly population 7,56 % and in a year with the 2011 69,65 % 7,58 elderly population .According to the distribution of the elderly.



Menopause is the end of the menstrual cycle routine permanently resulting from the loss of the activity of the ovary. Said to have a woman can menopause when the menstrual cycle berturut-turut stop for at least 12 months .This condition usually occurs in old lady 50-55 years .It s (Abramovitz 2020). Bleeding enough pascamenopause are complaints pascamenopause often experienced by women .Bleeding pascamenopause occurred at around 4-11 % woman is menopause .Although most cases are mild disease pascamenopause bleeding , 10 % cases caused by malignancy , the endometrial cancer .The highest insidensi endometrial cancer found in women aged 60 pascamenopause 79 years .It s (Abramovitz 2020).

2. METHOD

The kind of case study of writing is a case study by the observational descriptive with the approach the case study method research conducted by means of research the problem through a case consisting of a single unit (Notoatmodjo, 2010). This case study using the obstetrics care 7 varney from collecting a baseline data to evaluation and the development of using SOAP. The case study is the location where the case was conducted (Notoatmodjo , 2005). In the preparation of this case study authors take ngesrep locations in the. The case study subjects research that is a subject that intended for examination by researchers Arikunto (research or target ,2006) .Subjects used in this case is Mrs. I P2A0 with hemorrhage Ngesrep 2 months in public health. This case study implemented beginning on the 21 to July 22 until finish. Instrumen used writer for data collection is format for the assessment the care of obstetrics by using 7 step management obstetrics Varney .The technique of the collection of data using primary data from anamnesa and physical examination.

3. RESULT AND DISCUSSION

In this case the assessment is based on subjective data of the state of Mrs. I experiencing complaints of brownish red blood discharge 2 during the month and there is a sense of heat from the face spread throughout the body, and followed by the onset of redness in the face and neck and sweating at night since 1 Week ago, objective data on Mrs.I mom's condition is anxious. Blood

pressure : 140/90 mmhg, Pulse : 80x / min , Temperature : 38,5 C, Respiration : 20x / min.

Midwifery care in Mrs. I with postmenopausal bleeding monitor the general condition and vital signs, as well as provide counseling about the changes that occur during menopause, recommends to exercise, recommends to consume nutritious foods containing Vitamin B complex such as green vegetables and side dishes namely tofu and tempeh, consume foods containing vitamin E such as wheat, beans, eel, fish oil, egg yolks, beans, lettuce, broccoli and sweet potatoes provide drug therapy clonidine 0.1 2 x A Day 6 tablets, vitamin B 10 tablets 1 x a day and FE 1 x Day 10 tablets. Further, a referral to dr.SpOG for further inspection.

4. CONCLUSION

Assessment of subjective and objective data carefully degan able to establish the correct diagnosis. for further treatment in accordance with the needs of the patient. the patient is confirmed to have menopause and abnormal bleeding leads to the occurrence of a disease. in primary facilities, collaboration actions can be carried out to secondary facilities.

REFERENCES

- [1] Arikunto, S. (2006). Dasar-dasar evaluasi penelitian. Jakarta: Bumi Aksara.
- [2] Asih, D. A. (2017). Hubungan tingkat pengetahuan dengan tingkat kecemasan pada perempuan menopause di wilayah kerja puskesmas Ciputat Tangerang Selatan. (Tesis yang tidak diterbitkan). UIN Syarif Hidayatullah Jakarta: Fakultas Kedokteran dan Ilmu Kesehatan.
- [3] Azhari, T. R., & Mirza, M. (2016). Hubungan Regulasi Diri dengan Kecemasan Menghadapi Dunia Kerja pada Mahasiswa Tingkat Akhir Universitas Syiah Kuala. *Mediapsi*, 2(2), 23-29.
- [3] Brown, L., Bryant, C., Brown, V., Bei, B., & Judd, F. (2015). Investigating how menopausal factors and selfcompassion shape well-being: An exploratory path analysis. *Maturitas*, 81(2), 293-299.
- [4] Friedman, H. S., & Schustack, M. W. (2008). Kepribadian teori klasik dan riset modern. Jakarta: Erlangga.
- [5] Hawari, D. (2008). Manajemen stress, cemas dan depresi. Jakarta: Balai Penerbit



- [6] Indrias, H. D., & Maliya, A.(2015). Hubungan antara perubahan fisik dengan perubahan psikologis wanita pada masa menopause di Kelurahan Pucang
- [7] Kothiyal P, Monika S. 2013. Post Menopausal Quality of Life and Associated Factors-A Review, *Journal of Scientific and Innovative Research*, 2 (4): 814- 823.
- [8] Kuntjoro, Zainuddin Sri. 2014. Menopause. Diakses dari URL : http://www.psychoshare.com/file640/psikologi_dewasa/menopause.html
- [9] Kalarhoudi, M.A., Taebi, M., Sadat, Z., Saberi, F, 2011. Assessment of quality of life in menopausal periods : A population study in Kashan, Iran *Red Crescent Med J*. 2011; 13(11):811-817
- [10] Notoatmodjo, Soekidjo. 2018. Pendidikan Dan Perilaku Kesehatan. Jakarta : Indonesia Nurningsih. 2012. Hubungan Tingkat Pengetahuan tentang Menopause
- [11] Mulyani, N. S. (2013). Menopause akhir siklus menstruasi pada wanita di usia pertengahan. Yogyakarta: Nuha Medika.
- [12] Proverawati, A. (2017). Menopause dan Sindrome Premenopause. (W. Kristiyanasari, Ed.). Yogyakarta: Nuha Medika.
- Asriati, C. R., Wijaya, M., Nirmala, S. A., Gondodiputro, S., & Rahmiati, L. (2019).
- [13] Gambaran Pengetahuan Ibu tentang Persiapan Fisik dan Psikis Memasuki Masa Menopause. *Jurnal Kesehatan Vokasional*, 4(2), 99. <https://doi.org/10.22146/jkesvo.41638>
- [14] Indarwati, & Maryatun. (2019). Karakteristik Wanita Menopause dan Perubahan Pola Seksualitas di Desa Kedungan. *Jurnal GASTER*, 17(1), 1–16. <https://doi.org/https://doi.org/10.30787/gaster.v17i1.293>
- [15] Kemenkes 2018. Data Dan Informasi Profil Kesehatan Indonesia. Kesehatan, K., & Indonesia, R. (n.d.). profil-kesehatan-Indonesia-2015.
- [16] Sibagariang, E. E. (2016). Kesehatan Reproduksi Wanita. Jakarta Timur: CV Trans Info Media.
- [17] Archer, D. F., Sturdee, D. W., Baber, R., Villiers, T. J. De, Pines, A., Freedman, R.R., Warren, M. (2011). Menopausal hot flushes and night sweats : where are we now ?, 515–528. <https://doi.org/10.3109/13697137.2011.608596>
- Chou, C.-C. (2013). MEMORY PERFORMANCE AMONG TAIWANESE POSTMENOPAUSAL WOMEN.
- [18] Koeryaman, M. trisyani, & Ermiami. (2018). Adaptasi gejala menopause dan pemenuhan kebutuhan seksual wanita wanita usia 50 - 60 tahun, 16(1), 21–30.
- Kumalasari, & Andhyantoro. (2012). Kesehatan reproduksi. Jakarta: Salemba Medika.
- [19] Vidayati, L. A. (2017). ANALISIS HUBUNGAN PENGGUNAAN KONTRASEPSI PADA WANITA DENGAN USIA MENOPAUSE.
- [20] Purwati, A., & Ismiatun. (2018). Menarche, Kejadian Menopause, Ibu, 2(6).
- Rottie, J., Mongi, T. O., & Wae, A. (2019). HUBUNGAN JUMLAH PARITAS DENGAN USIA MENOPAUSE PADA WANITA DI DESA BULAN JAYA KABUPATEN TOJO UNA-UNA